


Kazan Golovkinsky Stratigraphic Meeting

2014



Kazan Federal University
Institute of Geology and Petroleum Technologies

Carboniferous and Permian Earth systems, stratigraphic events, biotic evolution, sedimentary basins and resources



Наземная растенія и плщеры
Сонсhifera

Вращіор о'д'и
Сонсhifera

(Наземная растенія и плщеры)?
Лингюта

Kazan Federal University
Institute of Geology and Petroleum Technologies

Kazan Golovkinsky Stratigraphic Meeting, 2014

Carboniferous and Permian Earth systems, stratigraphic events, biotic evolution, sedimentary basins and resources

October, 20–23, Kazan, Russia

Abstract Volume

Responsible editor
Danis K. Nurgaliev

Technical editors:
Vladimir V. Silantiev,
Milyausha N. Urazaeva

Reconstruction of sedimentary conditions of Middle Permian Kama-Ural basin studied by N.A.Golovkinsky

Alexander Lalomov¹, Guy Berthault², Marina Tugarova³, Viktor G. Isotov⁴, Lyalya M. Sitdikova⁴

¹ Institute of Geology of Ore Deposits, Petrography, Mineralogy and Geochemistry, Russian Academy of Science IGEM RAS, Moscow, Russia

² International Association of Sedimentologists, France

³ Saint-Petersburg State University, St-Petersburg, Russia

⁴ Kazan Federal University, Kazan, Russia

Golovkinsky Law (also known in Europe as Walter's Law) is one of the fundamental regularities that characterize lithological and facial features of sedimentary formations. The essence of the Law is successive alternation of types and character of sedimentation processes in sedimentary basins according to changes of hydrodynamic and hydrochemistry of the sedimentation environment both in a lateral and vertical direction. N. Golovkinsky provided the conclusions based upon research of a succession of sedimentary rocks in the Permian sequence of the valley of the Kama River [1].

Detail research of lithological structure, granulometric content and textural features of Middle Permian strata of Ural-Kama basin allows providing reconstruction of paleohydraulic conditions of sedimentation. Obtained data gave quantity illustration of Golovkinsky law of facial substitution.

Researched area mainly belongs to zone of development of sedimentation basin of Middle Permian age. Sustainable negative tectonic movements had beginning westward of Ural Folded System in Middle Permian and reached maximal expansion in Kazan time. Consecutive change of facies forms mainly terrigenous deposits in the east part and carbonate-clay, carbonate and carbonate-sulphate sequence in the west. In the zone of interrelations, it was complicated with shallow-sea lens-shape bodies of sandstones and conglomerates fixed geodynamically active structures of Tatar Arch, which divide the basin to terrigenous (east) and mostly clay-carbonate-sulphate (west) parts. The Kazan sequence consists of two series: transgressive Lower Kazan and regressive Upper Kazan [2].

Analysis of paleoflows based on research of structures and textures of Kazan stage sediments allows revealing of paleohydraulic conditions in the sedimentary basin.

Diagram of Rubin and McCulloch (1980) allows estimation of the flow velocity by parameters of cross-beds; the same data are possible to receive from granulometric composition using Hjulstrom diagram (Hjulstrom, 1935). The directions of the flows were obtained from rose-diagram of cross-beds.

Whereas in the continental pre-Ural part velocity of the flows orientated mainly from east to west reached 1.8 m/s in the beginning of transgression and decreased to 0.2-0.8 m/s in the end of regression stage, the situation in the marine part was much more complicated.

In the western (carbonate) basin the velocities varied from 0.2 m/s in the beginning of transgression to zero in the upper point.

In the area of Tatar Arch during Ufa Stage the paleoflows had south and south-east directions and were orientated parallel to the coastline of paleobasin. Paleoflows of north-west to south-west prevailed in the Early Kazan Age.

The structure of paleoflows cardinaly changed in Later Kazan Age. In the north part of Tatar Arch south-west directions of paleoflows had been prevailed. In the south part both south-east and south-west paleoflows are revealed, that is concerning with moderate development of Tatar Arch structures. Velocities of the currents in the Arch zone reached 1.8 v/s.

Facial interrelations in the zone of Tatar Arch and western basin were researched in detail by joint group of Kazan University, Saint-Petersburg University and IGEM RAS. In the westward direction facial transition from coarse terrigenous sediments (sandstones with pebbles) through siltstones and argillites to carbonate-gypsum sequence is observed. As the transgression developed, facial zones shifted after the coastline; therefore, coarser sediments had been overlay by finer ones.

For example, lower part of Early Kazan sequence near the axis of Tatar Arch (Elabuga – Krasny Bor) is represented by sandstones with pebbles of argillites up to 3-5 cm. Westward in the mouth of Vyatka the deposits are laterally changed into sandstone-siltstone-argillite strata. Far to the west the terrigenous deposits change to carbonates. At the same time, in vertical direction sandstones changed to siltstones, argillites – to carbonates and carbonates – to carbonate-gypsum lowers consequently. The transgression sequence in the facial interrelations area of Tatar Arch is complicated by consedimentation tectonic movements.

Thus, observed lithological lowers are not corresponds to a single time interval, but belongs to different sliding geochronological divisions. The sedimentary genesis of the sequence is explained by simultaneous prograding of the strata both laterally and vertically and movement of facial zoning during transgression and regression cycles.

This explanation does not corresponds to the classical Stenon stratigraphic concept of successive horizontal layers, but agree with Golovkinsky law which become foundation of sequence stratigraphy.

Golovkinsky N.A. (1868): About Permian formation in central part of Kama-Volga basin. *Proceedeengs of Geology of Russia* 1, 2: 146 p. (in Russian)

Geology of Tatarstan (2003): stratigraphy and tectonics. Moscow: GEOS: 402 p. (in Russian).

Rubin, D.M., McCulloch, D.S. (1980): Single and Superposed Bed Forms: A Synthesis of San Francisco Bay and Flume Observations. – *J. Sediment. Petrol.*, 26: 207–231.

Hjulstrom, F. (1935): The Morphological Activity of Rivers as Illustrated by River Fyris. – *Bull. Geol. Inst., Uppsala*, 25: 89–122.